Test Name	Domains	Scoring	Features	Disadvantages
Lymphedema Breast Cancer Questionnaire (LBCQ)*[18]	Self-reported symptoms in the last 30 days or year (swelling, tightness, heaviness, redness, infection)	Yes scored as 2, No scored as 1.	Open ended questions on symptom management	
Freiburg Life Quality Assessment-Lymphedema	Physical status, everyday life, social life, emotional wellbeing, treatment, satisfaction with life, profession and household	5-point Likert scale Higher scores indicate greater impairment	A short form was developed containing 33 questions[19,20]	Very long (92 questions)
Lymphedema Functioning, Disability and Health Questionnaire (Lymph-ICF)*[21]	Physical and mental functioning, household activities, mobility, life and social activities	0-100 scale	Helps distinguish between lymphedema and breast cancer treatment effects	
Lymphedema Life Impact Scale (LLIS)[22,23]	Physical, psychosocial, and functional concerns, and infection occurrence in the last year	5-point Likert scale Infection recurrence measures 0-≥4 Higher scores indicate greater impairment	Second version available	Does not correlate with edema severity
Lymphedema Symptom Intensity and Distress Survey (LSIDS)[24,25]	Soft tissue and neurological sensation, function, biobehavioral, sexuality, activity and resources	5-point Likert Higher scores indicate greater impairment	Different questionnaires available for Arm, Trunk and Head and Neck	
Lymphedema Quality of Life Inventory (LyQLI)[26]	Physical, psychosocial, practical concerns	4-point Likert Higher scores indicate greater impairment	Shorter version of the Swedish Lymphedema Quality of Life Inventory	
Lymphedema quality of life questionnaire (LYMQOL)[27]	Symptoms, body image/ appearance, function, mood	4-point Likert Higher scores indicate greater impairment	Separate arm and leg questionnaire Validated	Not designed for use in children Does not specifically address midline edema
Upper Limb Lymphedema 27 (ULL-27)*[28]	Physical, psychological, and social functioning	5-point Likert Higher scores indicate greater impairment	Sensitive to clinical changes	Addresses only upper limb lymphedema

**Breast Cancer Specific*